

# Spectacle Lenses With Aspherical Lenslets for Myopia Control vs Single-Vision Spectacle Lenses

## A Randomized Clinical Trial

Jinhua Bao, PhD; Yingying Huang, MD; Xue Li, PhD; Adeline Yang, MSc; Fengchao Zhou, BSc; Junqian Wu, BSc; Chu Wang, BSc; Yuhao Li, BSc; Ee Woon Lim, BSc; Daniel P. Spiegel, PhD; Björn Drobe, PhD; Hao Chen, MD, OD

**IMPORTANCE** Reducing myopia progression can reduce the risk of associated ocular pathologies.

**OBJECTIVE** To evaluate whether spectacle lenses with higher lenslet asphericity have a higher myopia control efficacy throughout 2 years.

**DESIGN, SETTING, AND PARTICIPANTS** This double-masked randomized clinical trial was conducted between July 2018 and October 2020 at the Eye Hospital of Wenzhou Medical University in Wenzhou, China. Children aged 8 to 13 years with a cycloplegic spherical equivalent refraction (SER) of  $-0.75$  D to  $-4.75$  D and astigmatism with less than  $-1.50$  D were recruited. A data and safety monitoring committee reviewed findings from a planned interim analysis in 2019.

**INTERVENTIONS** Participants were randomly assigned in a 1:1:1 ratio to receive spectacle lenses with highly aspherical lenslets (HAL), spectacle lenses with slightly aspherical lenslets (SAL), or single-vision spectacle lenses (SVL).

**MAIN OUTCOME AND MEASURES** Two-year changes in SER and axial length and their differences between groups.

**RESULTS** Of 157 participants who completed each visit (mean [SD] age, 10.4 [1.2] years), 54 were analyzed in the HAL group, 53 in the SAL group, and 50 in the SVL group. Mean (SE) 2-year myopia progression in the SVL group was 1.46 (0.09) D. Compared with SVL, the mean (SE) change in SER was less for HAL (by 0.80 [0.11] D) and SAL (by 0.42 [0.11] D;  $P \leq .001$ ). The mean (SE) increase in axial length was 0.69 (0.04) mm for SVL. Compared with SVL, increase in axial length was slowed by a mean (SE) of 0.35 (0.05) mm for HAL and 0.18 (0.05) mm for SAL ( $P \leq .001$ ). Compared with SVL, for children who wore HAL at least 12 hours every day, the mean (SE) change in SER was slowed by 0.99 (0.12) D, and increase in axial length slowed by 0.41 (0.05) mm.

**CONCLUSIONS AND RELEVANCE** In this study, HAL and SAL reduced the rate of myopia progression and axial elongation throughout 2 years, with higher efficacy for HAL. Longer wearing hours resulted in better myopia control efficacy for HAL.

**TRIAL REGISTRATION** Chinese Clinical Trial Registry Identifier: [ChiCTR1800017683](https://www.clinicaltrials.gov/ct2/show/study?term=ChiCTR1800017683)

*JAMA Ophthalmol.* 2022;140(5):472-478. doi:10.1001/jamaophthalmol.2022.0401  
Published online March 31, 2022.

- [+ Visual Abstract](#)
- [← Invited Commentary page 478](#)
- [+ Supplemental content](#)

**Author Affiliations:** Eye Hospital and School of Ophthalmology and Optometry, Wenzhou Medical University, Wenzhou, Zhejiang, China (Bao, Huang, X. Li, Zhou, Wu, Wang, Y. Li, Chen); National Clinical Research Center for Ocular Diseases, Wenzhou, Zhejiang, China (Bao, Huang, X. Li, Chen); Wenzhou Medical University-Essilor International Research Center (WEIRC), Wenzhou Medical University, Wenzhou, Zhejiang, China (Bao, Huang, X. Li, Yang, Lim, Spiegel, Drobe); R&D AMERA, Essilor International, Singapore, Singapore (Yang, Lim, Spiegel, Drobe).

**Corresponding Author:** Hao Chen, MD, OD, Eye Hospital and School of Ophthalmology and Optometry, Wenzhou Medical University, 270 Xueyuan Rd, Wenzhou 325027, China ([chenhao@mail.eye.ac.cn](mailto:chenhao@mail.eye.ac.cn)).

The projected 2050 prevalence rates of myopia and high myopia are alarming.<sup>1</sup> Myopia control interventions have been used for many years to reduce the severity of myopia and decrease the risk of associated ocular pathologies.<sup>2</sup> Increasing evidence suggests that specifically designed optical interventions such as spectacle lenses, soft contact lenses, and orthokeratology slow myopia progression in children.<sup>3-5</sup> The common features of these myopia optical interventions are to provide central correction for distance vision and correct peripheral retinal defocus or induce peripheral myopic retinal defocus simultaneously. Peripheral visual signals have been found to dominate central refractive development,<sup>6</sup> and the effect of peripheral myopic retinal defocus was found to provide myopia control signals.<sup>7-9</sup> Furthermore, several studies have shown a positive dose-response relationship between the efficacy of optical interventions and parameters such as addition power in multifocal soft contact lenses<sup>10</sup> and spectacle lenses<sup>11,12</sup> and in the wearing time of multifocal soft contact lenses.<sup>13</sup> This study evaluates novel spectacle lenses with aspherical lenslets and explores the effect of lenslet asphericity on myopia control efficacy. The results from the first-year interim analysis showed that spectacle lenses with highly aspherical lenslets (HAL) and spectacle lenses with slightly aspherical lenslets (SAL) were effective in slowing myopia progression. Moreover, a dose-dependent response was observed, as HAL had significantly better myopia control effect than SAL.<sup>14</sup>

We aimed to evaluate whether spectacle lenses with aspherical lenslets still slow myopia progression over 2 years and whether the level of lenslet asphericity would still affect myopia control efficacy in a dose-dependent manner.

## Methods

### Study Design

This 2-year, double-masked, 3-group, parallel randomized, single-center clinical trial comparing the effect of spectacle lenses with aspherical lenslets with single-vision spectacle lenses (SVL) in slowing myopia progression among children was conducted in the Eye Hospital of Wenzhou Medical University in Wenzhou, China, from July 18, 2018, to October 7, 2020. The study design has been described previously.<sup>14</sup> The trial protocol, standard operating procedures, and statistical analysis plan are available in [Supplement 1](#), [Supplement 2](#), and [Supplement 3](#), respectively. Children were randomly assigned to wear HAL, SAL, or SVL based on their baseline right eye cycloplegic spherical equivalent refraction (SER), age, and sex. Assignment with online software (Randola<sup>15</sup>) was performed by a specified person who had no interaction with the participants and stored on the website that required login credentials. The same person would pack the lenses according to the randomization in an unlabeled envelope with participant identification as the only identification for dispensing. Masking was maintained for refraction and axial length measurements for dispensing personnel and examiners, participants, and parents or guardians. However, it was not entirely possible to mask HAL and SAL lenses from SVL because lenslets were visible un-

### Key Points

**Question** What is the efficacy of spectacle lenses with highly aspherical lenslets and slightly aspherical lenslets compared with conventional single-vision spectacle lenses in controlling myopia progression throughout 2 years?

**Findings** In this randomized clinical trial of 157 children, the amount of myopia progression and axial length increase was significantly less in both the highly aspherical lenslets group and the slightly aspherical lenslets group vs the single-vision spectacle lenses group at 24 months.

**Meaning** The findings of this study suggest that a higher asphericity of lenslets may be associated with more effective myopia control.

der certain lighting conditions. However, complete masking was possible between HAL and SAL. Written informed consent was obtained from the parent or guardian, and written assent was obtained from the study participant. All study participants were given a pair of myopia-control spectacles on completion of the study. A data and safety monitoring committee reviewed the trial data for participant safety. The ethics committee of Eye Hospital of Wenzhou Medical University approved the clinical trial on July 17, 2018, and the trial was registered with the Chinese Clinical Trial Registry. All procedures adhered to the tenets of the Declaration of Helsinki.<sup>16</sup>

### Participants

The first participant was enrolled on August 11, 2018. Chinese children aged 8 to 13 years who had myopia, SER from  $-0.75$  diopters (D) to  $-4.75$  D, astigmatism of 1.50 D cylinder or less, anisometropia of 1.00 D or less, and best-corrected visual acuity of 0.05 logMAR or better in each eye were randomized at a 1:1:1 ratio to wear HAL, SAL, or SVL. No participants had a history of myopia control or any ocular or systemic issues that could affect vision outcomes.

### Study Procedures


A detailed description of this study device can be found elsewhere.<sup>14</sup> The primary outcomes were changes in SER and axial length. The parameter measurement procedures followed those for the 1-year outcomes of HAL and SAL.<sup>14</sup> SER and axial length were measured every 6 months. SER was measured by the mode of 10 measurements using a Topcon KR-800 (Topcon Corporation), which was acquired at least 30 minutes after installing 2 drops of cyclopentolate, 1%, administered 5 minutes apart. Axial length was calculated as the mean of 5 measurements obtained using a Lenstar LS 900 instrument (Haag-Streit AG).

The secondary outcomes presented in this report were lens wearing hours assessed by 6-month questionnaires with self-reported wearing times per day, every day per week (Monday to Sunday).

### Statistical Analysis

Based on previous findings,<sup>12,17,18</sup> we anticipated a mean (SD) SER progression of 1.50 (0.75) D and converted axial length progression of 0.6 (0.02) mm in the control group throughout a

Figure 1. CONSORT Study Flowchart



2-year period. We wanted to identify a 33% reduction in the amount of SER and axial length progression for treatment groups compared with the control group. Fifty participants were required in each group for an  $\alpha$  level of .05 (2-tailed), a power of 90%, and a dropout rate of 10%. With an interim analysis at 1 year, the  $\alpha$  level was adjusted to .029 based on the Pocock method for 2-year outcomes of SER and axial length.

All data from participants who had complete 2-year follow-up records were analyzed. The mean values for ocular parameters measured in the right eye were used because there was a high correlation between the 2 eyes in SER ( $r = 0.74$ ,  $P < .001$ ) and axial length ( $r = 0.87$ ,  $P < .001$ ). The changes in SER and axial length from baseline are presented as mean (SE).  $\chi^2$  Test and analysis of variance with post hoc Bonferroni test were used to assess intergroup differences in categorical and continuous variables, respectively. Our analysis was performed using complete case data without imputation for missing data and dropouts. We undertook analyses using linear mixed model, adjusted for baseline age, sex, SER, axial length, age at myopia onset, and number of parents with myopia to evaluate the treatment effect. The mean daily lens wearing hours were calculated based on the mean daily lens wearing hours over the four 6-month periods. The effect of wearing time was analyzed categorically. Full-time wearers were defined as children who reported wearing their study devices for at least 12 hours every day. Part-time wearers were defined as nonfull-time wearers. SPSS statistical software version 24 (IBM) was used for data analysis. Two-sided  $P$  values of less than .05 were considered statistically significant.

## Results

One hundred seventy children with a mean (SD) age of 10.4 (1.2) years, ranging from 8 to 13 years, were recruited and randomized among the HAL ( $n = 58$ ), SAL ( $n = 57$ ), and SVL ( $n = 55$ ) groups (Figure 1). Only 167 were dispensed with the study equipment. Three children discontinued the study: 1 pre-

sented with intermittent exotropia not apparent during screening, 1 belatedly reported history of using myopia control, and 1 dropped out before test lenses were dispensed. After 2 years, 157 participants had completed all their visits; among children who did not attend their follow-up appointments, 2 (3.6%), 3 (5.4%), and 4 (7.3%) were in the HAL, SAL, and SVL groups, respectively. The reasons for dropout were not related to study devices. The demographic and ocular characteristics of each group at baseline are shown in Table 1.

### Primary Outcome

#### Changes in Spherical Equivalent Over 2 Years

The 2-year means (SEs) for myopia progression were  $-0.66$  (0.09) D,  $-1.04$  (0.06) D, and  $-1.46$  (0.09) D in the HAL, SAL, and SVL groups, respectively (Figure 2). Significant differences were found among treatment groups ( $F_{2,154} = 25.80$ ;  $P < .001$ ). The HAL and SAL groups had less SER progression by a mean (SE) of 0.80 (0.11) D (95% CI, 0.53-1.07;  $P < .001$ ) and 0.42 (0.11) D (95% CI, 0.15-0.70;  $P = .001$ ), respectively, than the SVL group. In addition, the HAL group had less SER progression than the SAL group, with a mean (SE) difference of 0.38 (0.11) D (95% CI, 0.11-0.64;  $P = .002$ ).

In the linear mixed model analysis, baseline age (95% CI, 0.004-0.128;  $P = .04$ ) was directly, significantly associated with SER progression. The model-adjusted mean (SE) changes in SER were  $-0.68$  (0.07) D,  $-1.04$  (0.08) D, and  $-1.45$  (0.08) D for HAL, SAL, and SVL, respectively, with a significant effect of lens design ( $F_{6,154} = 7.88$ ;  $P < .001$ ) on SER. Compared with SVL, the adjusted differences in mean (SE) SER were 0.77 (0.11) D (95% CI, 0.51-1.04;  $P < .001$ ) and 0.42 (0.11) D (95% CI, 0.15-0.68;  $P = .001$ ) in the HAL and SAL groups, respectively, and 0.36 (0.11) D (95% CI, 0.10-0.62;  $P = .003$ ) between the HAL and SAL groups.


#### Changes in Axial Length Over 2 Years

The mean (SE) increases in axial length over 2 years were 0.34 (0.03) mm, 0.51 (0.04) mm, and 0.69 (0.04) mm in the HAL, SAL, and SVL groups, respectively (Figure 2). As with SER, there

**Table 1. Baseline Demographic and Ocular Characteristics of Participants Who Completed the 24-Month Follow-up in Each Group**

Clinical value	Mean (SE)		
	HAL (n = 54)	SAL (n = 53)	SVL (n = 50)
Age, y	10.6 (0.2)	10.2 (0.2)	10.4 (0.2)
Female, No. (%)	28 (52)	36 (68)	21 (42)
Male, No. (%)	26 (48)	17 (32)	29 (58)
Right eye			
Refractive error (SER), D	-2.70 (0.14)	-2.28 (0.13)	-2.44 (0.12)
Axial length, mm	24.76 (0.09)	24.44 (0.10)	24.77 (0.09)
Age at myopia onset, y	9.3 (0.2)	9.4 (0.2)	9.3 (0.2)
Parents with myopia, No. (%)			
0	18 (33.3)	12 (22.6)	12 (24.0)
1	20 (37.0)	22 (41.5)	18 (36.0)
2	16 (29.6)	19 (35.8)	20 (40.0)

Abbreviations: D, diopters; HAL, spectacle lenses with highly aspherical lenslets; SAL, spectacle lenses with slightly aspherical lenslets; SER, spherical equivalent refraction; SVL, single-vision spectacle lenses.

**Figure 2. Change in Unadjusted Spherical Equivalent Refraction (SER) and Axial Length (AL) From Baseline to 2 Years**

Error bars represent standard errors of the mean. HAL indicates spectacle lenses with highly aspherical lenslets; SAL, spectacle lenses with slightly aspherical lenslets; SVL, single-vision spectacle lenses.

was a significant difference ( $F_{2,154} = 24.98$ ;  $P < .001$ ) in axial length increase among treatment groups. Compared with the SVL group, the HAL and SAL groups had reduced mean (SE) axial length elongation by 0.35 (0.05) mm (95% CI, 0.23-0.47;  $P < .001$ ) and 0.18 (0.05) mm (95% CI, 0.06-0.30;  $P = .001$ ), respectively. Furthermore, HAL had less axial length elongation than SAL by a mean (SE) of 0.17 (0.05) mm (95% CI, 0.05-0.29;  $P = .002$ ).

In the linear mixed model analysis, baseline age (95% CI, -0.04 to -0.01;  $P = .002$ ) and age at myopia onset (95% CI, 0.003-0.032;  $P = .02$ ) were significantly associated with axial length elongation. After adjustment, the means (SEs) for change in axial length were 0.35 (0.03) mm, 0.50 (0.03) mm, and 0.68 (0.03) mm for HAL, SAL, and SVL, respectively, with 0.34 (0.05) mm (95% CI, 0.23-0.45;  $P < .001$ ) and 0.18 (0.05) mm (95% CI, 0.07-0.30;  $P < .001$ ) reductions in axial length elongation in the HAL and SAL groups compared with SVL. The HAL group had less axial length increase than SAL by a mean (SE) of 0.16 (0.05) mm (95% CI, 0.05-0.27;  $P = .002$ ). As with SER, there was a significant effect of lens design ( $F_{6,154} = 8.41$ ;

$P < .001$ ) in axial length increase among treatment groups throughout 2 years.

### Secondary Outcomes

During the second year, HAL still slowed myopia progression compared with SVL (mean [SE] for HAL, SER: -0.39 [0.05] D;  $P < .001$ ; axial length, 0.21 [0.02] mm;  $P < .001$ ). However, there were no differences in SER and axial length change observed between the SAL and SVL groups during the second year (mean [SE] for SER, SAL: -0.57 [0.05] D; SVL: -0.64 [0.05] D;  $P = .90$ ; axial length, SAL: 0.27 [0.02] mm; SVL: 0.32 [0.02] mm;  $P = .12$ ). Thus, compared with SVL, SAL slowed myopia progression of children mainly during the first year.

The mean daily wearing time throughout 2 years was similar among the groups, with mean (SE) durations of 13.4 (0.29) hours, 13.4 (0.24) hours, and 13.9 (0.24) hours for HAL, SAL, and SVL, respectively ( $F_{2,154} = 1.08$ ;  $P = .34$ ). The mean wearing hours increased throughout the study from 13.1 hours in the first year to 14.0 hours in the second year (95% CI, 0.47-1.17;  $P < .001$ ), with 61% ( $n = 96$ ) of children who were full-

Table 2. Comparisons of Myopia Progression and Axial Elongation Between Full- and Part-time Wearers

Clinical value	Mean (SE)		
	HAL <sup>a</sup>	SAL <sup>a</sup>	SVL <sup>a</sup>
<b>Change in SER, D</b>			
Full-time wearers	-0.48 (0.10)	-0.95 (0.08)	-1.44 (0.10)
Part-time wearers	-0.93 (0.13)	-1.15 (0.10)	-1.50 (0.16)
P value	.01	.11	.77
<b>Change in AL, mm</b>			
Full-time wearers	0.28 (0.04)	0.46 (0.04)	0.69 (0.04)
Part-time wearers	0.43 (0.06)	0.57 (0.04)	0.70 (0.07)
P value	.03	.10	.92

Abbreviations: AL, axial length; HAL, spectacle lenses with highly aspherical lenslets; SAL, spectacle lenses with slightly aspherical lenslets; SER, spherical equivalent refraction; SVL, single-vision spectacle lenses.

<sup>a</sup> Full-time wearers wore lenses at least 12 hours per day, and part-time wearers

wore lenses less than 12 hours per day. In the HAL group, there were 32 full-time wearers and 22 part-time wearers. In the SAL group, there were 30 full-time wearers and 23 part-time wearers. In the SVL group, there were 34 full-time wearers and 16 part-time wearers.

time wearers in the first year to 89% (n = 139) who were full-time wearers in the second year ( $P < .001$ ). Therefore, we analyzed the myopia control efficacy based on the first-year full and part-time categories. The SER progression in the SAL and SVL groups was similar for full-time wearers and part-time wearers but was lower for full-time wearers in the HAL group (mean [SE] difference, 0.45 [0.16] D;  $P = .01$ ; Table 2). Axial length elongation for full-time wearers was similar to that among part-time wearers in the SAL and SVL groups but was lower for full-time wearers in the HAL group (mean [SE] difference, 0.15 [0.07] mm;  $P = .03$ ; Table 2). For full-time HAL wearers, the mean (SE) 2-year treatment effect compared with SVL group was 0.99 (0.12) D for SER and 0.41 (0.05) mm for axial length, whereas for HAL part-time wearers, it was 0.54 (0.15) D for SER and 0.26 (0.07) mm for axial length.

### Adverse Events

None of the adverse events reported were severe or resulted in study discontinuation. Two events were mild and unrelated to the study device and resolved with no reported loss of best-corrected visual acuity. One SVL wearer had punctate corneal epithelial defects, and the other SAL wearer was hit by a ball, causing the spectacle frame to break and the participant's face to be bruised. There was no significant difference in adverse events between treatment groups ( $P = .59$ ).

### Discussion

In this 2-year randomized clinical trial, HAL slowed myopia progression by 0.80 D (55%) and increase in axial length by 0.35 mm (51%) compared with SVL. Compared with SAL, HAL slowed myopia progression by 0.38 D (37%) and axial length by 0.17 mm (33%). This outcome is well in line with the first-year interim analysis,<sup>14</sup> and it confirms a positive dose-response relationship between myopia control efficacy and lenslet asphericity. The dose-dependent effect of optical interventions in minimizing lens-induced myopia in animal studies was attributed mainly to lens design features such as the amount and area of lens addition,<sup>19,20</sup> peripheral defocus, and lens asphericity.<sup>21,22</sup> This dose-dependent relationship was also found in human clinical

studies for ophthalmic lenses with higher addition<sup>10,12</sup> and contact lenses with higher asphericity.<sup>23</sup> Compared with SVL, during the second year, HAL remained effective in slowing myopia progression. This was not the case for SAL that slowed myopia progression of children mainly during the first year. This is similar to the previous myopia control study using progressive addition lenses.<sup>24</sup> It may be speculated that there is also a dose-response relationship between the duration of myopia control efficacy and lenslet asphericity. Therefore, it will be interesting to evaluate whether myopia control efficacy of HAL extends beyond 2 years.

Bifocal, prismatic bifocal spectacle lenses and defocus incorporated multiple segments (DIMS) spectacle lenses have also shown clinically significant myopia control results in Chinese children over the same duration as our study.<sup>25,26</sup> Throughout 2 years, the myopia control efficacy for HAL (0.80 D) and prismatic bifocal lenses (0.85 D) were comparable, while SAL (0.42 D), bifocal lenses (0.59 D), and DIMS (0.44 D) had lower myopia control efficacy. In terms of reducing axial elongation, HAL (0.35 mm) and DIMS (0.34 mm) had higher efficacy than SAL (0.18 mm) and bifocal and prismatic bifocal lenses (0.21 mm). Differences in myopia control efficacy could be linked to differences in lens designs, namely, the concentric ring configuration with aspherical lenslets (this study), honeycomb configuration with spherical lenslets (DIMS),<sup>27</sup> or increase in power over the lower part of the lens (bifocal and prismatic bifocal lenses). Another reason might be related to the control groups. Myopia progression rates among SVL wearers were -1.46 D in the present study and -1.55 D in the study by Cheng et al,<sup>25</sup> which are consistent with previous findings in Chinese children with myopia<sup>28,29</sup> while in the DIMS study, children wearing SVL progressed significantly less (-0.85 D).

The effect of wearing time on myopia control efficacy of HAL also indicated a similar dose-response relationship, increasing myopia control efficacy to 0.99 D (67%) and 0.41 mm (60%) for full-time wearers (at least 12 hours per day). A better myopia control effect with increased wearing time was also found in a study using defocus incorporated soft contact lenses.<sup>13</sup> However, no correlation between lens wearing time and myopia control efficacy was found in the study using DIMS spectacle lenses.<sup>26</sup> In the DIMS study, participants wore their

study devices constantly for more than 15 hours per day.<sup>26</sup> In contrast, 39% of participants in this study wore their spectacles for less than 12 hours every day in the first year and 11% in the second year. Throughout 2 years, myopia progression and axial elongation were significantly lower among full-time wearers than among part-time wearers in the HAL group but not in the SVL or SAL group. As such, full-time wearing of myopia control intervention should be recommended for better outcome.

### Strengths and Limitations

The first notable strength of this study is its low dropout rate of 6.5%. Second, the recruitment duration was short (approximately 2 months), with very punctual follow-ups (except the 18-month visit). This minimized the possible influence of confounding factors in the study, such as seasonal effects on myopia progression.<sup>30</sup>

Nevertheless, there were some limitations to this study. First, this study was conducted with a strictly Chinese cohort in Wenzhou, China. According to the myopia screening of school-aged children (7 to 18 years) from more than 1000 elementary and high schools of Wenzhou in 2020, the overall myopia prevalence and high myopia prevalence were 59.35% and 4.99%, respectively,<sup>31</sup> which was comparable with the findings of 2 recent studies in China.<sup>32,33</sup> The current study sample

could represent school-aged children in China but does not necessarily represent children with myopia globally. Second, the precision of the questionnaire may not reflect actual wearing hours. An objective measure of wearing time would be preferred. Third, the 18-month visit was delayed by a mean of 3 weeks. As a result of COVID-19 causing an unprecedented global pandemic, a lockdown was imposed for 3 weeks in February 2020. Children did not physically attend school for 4 months from February to the end of May 2020.<sup>34</sup> Other than the delay during the 18-month visit, COVID-19 did not have any major impact on the conduct of the study. The 24-month visit was conducted according to the investigation plan. Lastly, outdoor time was not measured and it may affect myopia progression in the study.

### Conclusions

In children with myopia, wearing HAL significantly reduced the rate of myopia progression and eye growth over 2 years compared with SAL and SVL. This study demonstrated a dose-dependent effect, with higher lenslet asphericity having greater myopia control efficacy. The full-time wearing of HAL increased myopia control efficacy to 0.99 D (67%) for SER and 0.41 mm (60%) for axial length.

#### ARTICLE INFORMATION

**Accepted for Publication:** February 1, 2022.

**Published Online:** March 31, 2022.  
doi:10.1001/jamaophthalmol.2022.0401

**Open Access:** This is an open access article distributed under the terms of the [CC-BY License](#). © 2022 Bao J et al. *JAMA Ophthalmology*.

**Author Contributions:** Dr Chen had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

**Concept and design:** Bao, Yang, Spiegel, Drobe, Chen.

**Acquisition, analysis, or interpretation of data:** All authors.

**Drafting of the manuscript:** Bao, Yang, Zhou, Wu, Wang, Y. Li, Lim, Spiegel, Drobe.

**Critical revision of the manuscript for important intellectual content:** Bao, Huang, X. Li, Yang, Spiegel, Drobe, Chen.

**Statistical analysis:** Bao, Huang, X. Li, Yang, Spiegel, Drobe.

**Obtained funding:** Bao, Drobe, Chen.

**Administrative, technical, or material support:** Yang, Lim.

**Supervision:** Bao, Drobe, Chen.

**Conflict of Interest Disclosures:** Dr Drobe reported pending patents for WO2019166653, WO2019166654, and WO2019166655 (co-inventor). Drs Yang, Spiegel, and Drobe and Ms Lim are employees of Essilor International, which sells lenses with highly aspherical lenslets design and supplied the study device. No other disclosures were reported.

**Funding/Support:** This study was supported in part by the International S&T Cooperation Program of China (grant 2014DFA30940) and the collaborative research project with Essilor

International (Wenzhou Medical University grants 95013006, 95016010, and 95020005).

**Role of the Funder/Sponsor:** Essilor International participated in the study design, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Data Sharing Statement:** See [Supplement 4](#).

#### REFERENCES

- Holden BA, Fricke TR, Wilson DA, et al. Global prevalence of myopia and high myopia and temporal trends from 2000 through 2050. *Ophthalmology*. 2016;123(5):1036-1042. doi:10.1016/j.ophtha.2016.01.006
- Saw SM, Gazzard G, Shih-Yen EC, Chua WH. Myopia and associated pathological complications. *Ophthalmic Physiol Opt*. 2005;25(5):381-391. doi:10.1111/j.1475-1313.2005.00298.x
- Wildsoet CF, Chia A, Cho P, et al. IMI: Interventions Myopia Institute: interventions for controlling myopia onset and progression report. *Invest Ophthalmol Vis Sci*. 2019;60(3):M106-M131. doi:10.1167/iov.18-25958
- Sankaridurg P, Conrad F, Tran H, Zhu J. Controlling progression of myopia: optical and pharmaceutical strategies. *Asia Pac J Ophthalmol (Phila)*. 2018;7(6):405-414. doi:10.22608/APO.2018333
- Huang J, Wen D, Wang Q, et al. Efficacy comparison of 16 interventions for myopia control in children: a network meta-analysis. *Ophthalmology*. 2016;123(4):697-708. doi:10.1016/j.ophtha.2015.11.010
- Smith EL III, Kee CS, Ramamirtham R, Qiao-Grider Y, Hung LF. Peripheral vision can influence eye growth and refractive development in infant monkeys. *Invest Ophthalmol Vis Sci*. 2005;46(11):3965-3972. doi:10.1167/iov.05-0445
- Smith EL III, Hung LF. The role of optical defocus in regulating refractive development in infant monkeys. *Vision Res*. 1999;39(8):1415-1435. doi:10.1016/S0042-6989(98)00229-6
- Wallman J, Gottlieb MD, Rajaram V, Fugate-Wentzek LA. Local retinal regions control local eye growth and myopia. *Science*. 1987;237(4810):73-77. doi:10.1126/science.3603011
- Smith EL III. Prentice Award Lecture 2010: a case for peripheral optical treatment strategies for myopia. *Optom Vis Sci*. 2011;88(9):1029-1044. doi:10.1097/OPX.0b013e3182279cfa
- Walline JJ, Walker MK, Mutti DO, et al; BLINK Study Group. Effect of high add power, medium add power, or single-vision contact lenses on myopia progression in children: the BLINK Randomized Clinical Trial. *JAMA*. 2020;324(6):571-580. doi:10.1001/jama.2020.10834
- Leung JT, Brown B. Progression of myopia in Hong Kong Chinese schoolchildren is slowed by wearing progressive lenses. *Optom Vis Sci*. 1999;76(6):346-354. doi:10.1097/00006324-199906000-00013
- Hasebe S, Jun J, Varnas SR. Myopia control with positively aspherized progressive addition lenses: a 2-year, multicenter, randomized, controlled trial. *Invest Ophthalmol Vis Sci*. 2014;55(11):7177-7188. doi:10.1167/iov.12-11462
- Lam CSY, Tang WC, Tse DYY, Tang YY, To CH. Defocus Inc Soft Contact (DISC) lens slows myopia progression in Hong Kong Chinese schoolchildren: a 2-year randomised clinical trial. *Br J Ophthalmol*. 2014;98(1):40-45. doi:10.1136/bjophthalmol-2013-303914

14. Bao J, Yang A, Huang Y, et al. One-year myopia control efficacy of spectacle lenses with aspherical lenslets. *Br J Ophthalmol*. 2021; [bjophthalmol-2020-318367](https://doi.org/10.1136/bjophthalmol-2020-318367). doi:10.1136/bjophthalmol-2020-318367
15. Randola. Accessed March 1, 2022. <http://www.rando.la/>
16. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;310(20):2191-2194. doi:10.1001/jama.2013.281053
17. Yang Z, Lan W, Ge J, et al. The effectiveness of progressive addition lenses on the progression of myopia in Chinese children. *Ophthalmic Physiol Opt*. 2009;29(1):41-48. doi:10.1111/j.1475-1313.2008.00608.x
18. Olsen T. Sources of error in intraocular lens power calculation. *J Cataract Refract Surg*. 1992;18(2):125-129. doi:10.1016/S0886-3350(13)80917-0
19. McFadden SA, Tse DY, Bowrey HE, et al. Integration of defocus by dual power Fresnel lenses inhibits myopia in the mammalian eye. *Invest Ophthalmol Vis Sci*. 2014;55(2):908-917. doi:10.1167/iov.13-11724
20. Tse DY, To CH. Graded competing regional myopic and hyperopic defocus produce summated emmetropization set points in chick. *Invest Ophthalmol Vis Sci*. 2011;52(11):8056-8062. doi:10.1167/iov.10-5207
21. Woods J, Guthrie SE, Keir N, et al. Inhibition of defocus-induced myopia in chickens. *Invest Ophthalmol Vis Sci*. 2013;54(4):2662-2668. doi:10.1167/iov.12-10742
22. Irving EL, Yakobchuk-Stanger C. Myopia progression control lens reverses induced myopia in chicks. *Ophthalmic Physiol Opt*. 2017;37(5):576-584. doi:10.1111/opo.12400
23. Cooper J, O'Connor B, Watanabe R, et al. Case series analysis of myopic progression control with a unique extended depth of focus multifocal contact lens. *Eye Contact Lens*. 2018;44(5):e16-e24. doi:10.1097/IJCL.0000000000000440
24. Brennan NA, Toubouti YM, Cheng X, Bullimore MA. Efficacy in myopia control. *Prog Retin Eye Res*. 2021;83:100923. doi:10.1016/j.preteyeres.2020.100923
25. Cheng D, Schmid KL, Woo GC, Drobe B. Randomized trial of effect of bifocal and prismatic bifocal spectacles on myopic progression: two-year results. *Arch Ophthalmol*. 2010;128(1):12-19. doi:10.1001/archophthalmol.2009.332
26. Lam CSY, Tang WC, Tse DY, et al. Defocus Inc Multiple Segments (DIMS) spectacle lenses slow myopia progression: a 2-year randomised clinical trial. *Br J Ophthalmol*. 2020;104(3):363-368. doi:10.1136/bjophthalmol-2018-313739
27. Li X, Ding C, Li Y, et al. Influence of lenslet configuration on short-term visual performance in myopia control spectacle lenses. *Front Neurosci*. 2021;15:667329. doi:10.3389/fnins.2021.667329
28. Yam JC, Jiang Y, Tang SM, et al. Low-concentration atropine for myopia progression (LAMP) study: a randomized, double-blinded, placebo-controlled trial of 0.05%, 0.025%, and 0.01% atropine eye drops in myopia control. *Ophthalmology*. 2019;126(1):113-124. doi:10.1016/j.ophtha.2018.05.029
29. Zhu MJ, Feng HY, He XG, Zou HD, Zhu JF. The control effect of orthokeratology on axial length elongation in Chinese children with myopia. *BMC Ophthalmol*. 2014;14(1):141. doi:10.1186/1471-2415-14-141
30. Fulk GW, Cyert LA, Parker DA. Seasonal variation in myopia progression and ocular elongation. *Optom Vis Sci*. 2002;79(1):46-51. doi:10.1097/00006324-200201000-00012
31. Xu L, Ma Y, Yuan J, et al. COVID-19 quarantine reveals that behavioral changes have an effect on myopia progression. *Ophthalmology*. 2021;127(11):1652-1654. doi:10.1016/j.ophtha.2021.04.001
32. Wang J, Li Y, Musch DC, et al. Progression of myopia in school-aged children after COVID-19 home confinement. *JAMA Ophthalmol*. 2021;139(3):293-300. doi:10.1001/jamaophthalmol.2020.6239
33. Chang P, Zhang B, Lin L, et al. Comparison of myopic progression before, during, and after COVID-19 lockdown. *Ophthalmology*. 2021;128(11):1655-1657. doi:10.1016/j.ophtha.2021.03.029
34. Zhang H, Guo LW, Gao YY, Yao H, Xie ZK, Zhang WX. The impact of the COVID-19 pandemic on pediatric clinical practice in Wenzhou, China: a retrospective study. *Front Pediatr*. 2020;8:585629. doi:10.3389/fped.2020.585629

## Invited Commentary

## Clinical Relevance of Myopia Control With Specialized Spectacles

David C. Musch, PhD, MPH; Steven M. Archer, MD

Based on a randomized clinical trial of spectacle lenses with aspherical lenslets that induce either slight or high asphericity, Bao et al<sup>1</sup> report the 2-year findings, which lend additional credence to the interim, 1-year findings.<sup>2</sup> Children aged 8 to 14 years with mild to moderate myopia (−0.75-diopters [D] to −4.75-D spherical equivalent refraction) randomly assigned to wear slightly aspherical lenslet (SAL) or highly aspherical lenslet (HAL) spectacles showed less myopic progression than those who wore single-vision spectacle lenses. In addition, the HAL wearers showed less progression than the SAL wearers.

The authors powered their study to detect a 33% reduction in the amount of spherical equivalent refraction and axial length progression, and they enrolled 170 children, 20 more than the 150 (50 per group) required to provide adequate power. They analyzed the 2-year, right-eye findings of 157 children (92.4%) and describe the reasons for lack of follow-up well. After 2 years of follow-up, children who wore the HAL or SAL spectacles had 0.80 D and 0.42 D less myopic progression, respectively, than wearers of single-vision spectacle lenses, with corresponding axial length findings.

Although the authors only analyzed information from the right eye—thus sacrificing the additional statistical power of

using both eyes with appropriate treatment of the intereye correlation<sup>3</sup>—the statistical significance of their findings is clear. What remains to be considered is the clinical relevance of their findings and the feasibility of having children wear spectacles with aspherical lenslets for many years. A recent study<sup>4</sup> that evaluated the benefits of each additional 1 D of myopic control on adverse outcomes places a value on the amount of myopic control that Bao et al<sup>1</sup> report. Given the children's mean (SD) age at enrollment of 10.6 (0.2) years, with much of their period of rapid myopic progression behind them, one wonders how much additional benefit would occur beyond 2 years. Would initiating aspheric spectacles earlier in life potentially provide more dramatic myopia control? Further study involving younger children is warranted.

However, would younger children adhere well to wearing spectacles with glare from the edges of the lenslets and degraded vision through the aspheric portion of the lens when the child's gaze wanders away from primary position? Wear time in this study was moderately worse for aspheric lenses, with approximately 58% of the HAL and SAL groups vs 68% of the single-vision spectacle lens group having worn their spectacles for at least 12 hours per day. The fact that the mean daily wear time increased from the first to second year in the HAL and SAL groups suggests that the children adjusted well to